

Clayton Metropolis

An oil dispersion formulation containing 10 g/L mesosulfuron-methyl and 2 g/L iodosulfuron-methyl-sodium. A highly active herbicide (a combination of two sulfonylurea herbicides) with foliar and some root activity against black-grass, wild oats, ryegrasses, meadow-grasses, common chickweed and mayweeds in winter wheat.
MAPP 18401

 	<p>CLAYTON METROPOLIS : An oil dispersion formulation containing 10 g/L mesosulfuron-methyl and 2 g/L iodosulfuron-methylsodium. Also contains mefenpyr-diethyl, solvent naphtha (petroleum), heavy aromatic and solvent naphtha (petroleum), light aromatic.</p> <p>Signal Word : WARNING Causes serious eye irritation. Very toxic to aquatic life with long lasting effects. Wear protective gloves/protective clothing/eye protection/face protection. If eye irritation persists: Get medical advice/ attention.</p> <p>Dispose of contents/container to a licensed hazardous-waste disposal contractor or collection site except for empty clean containers which can be disposed of as non-hazardous waste. Contains fatty alcohol ethoxylate alkyl ether. May produce an allergic reaction.</p>
<p>To avoid risks to human health and the environment, comply with the instructions for use.</p> <p>THE (COSHH) CONTROL OF SUBSTANCES HAZARDOUS TO HEALTH REGULATIONS MAY APPLY TO THE USE OF THIS PRODUCT AT WORK</p>	

IMPORTANT INFORMATION : FOR USE ONLY AS AN AGRICULTURAL HERBICIDE	
Crops :	Winter wheat
Maximum individual dose :	1.2 L product/ha
Maximum number of treatments :	One per crop
Latest time of application :	Flag leaf ligule just visible (GS 39) of the crop
<p>READ THE LABEL BEFORE USE. USING THIS PRODUCT IN A MANNER THAT IS INCONSISTENT WITH THE LABEL MAY BE AN OFFENCE. FOLLOW THE CODE OF PRACTICE FOR USING PLANT PROTECTION PRODUCTS.</p>	

<p>Approval Holder Clayton Plant Protection Ltd., Unit F10, Bracetown Business Park Clonee, Co. Meath. Ireland Tel: (00 353) 1 8210127 www.cpp.ag Email: info@cpp.ag</p> <p>Marketing company in UK Clayton Plant Protection (UK) Ltd. Address and telephone as above.</p>	<p>Contents: 5 Litres e</p> <p>SHAKE THOROUGHLY BEFORE USE</p> <p>PROTECT FROM FROST STORE IN A COOL DRY PLACE</p> <p>UN3082</p> <p>Batch No:</p>
--	---

Conditions of Supply: all goods supplied by us are of high quality and we believe them to be correct but, as we cannot exercise control over their storage, handling, mixing or use, or weather conditions before, during and after application which may affect the performance of the goods, all conditions and warranties, statutory or otherwise, as to the quality or fitness for any purpose of our goods are excluded, and no responsibility will be accepted by us or resellers for any failure in performance, damage or injury whatsoever arising from their storage, handling, application or use. These conditions cannot be varied by our staff or agents whether or not they supervise or assist in the use of such goods.

SAFETY PRECAUTIONS

Operator Protection

Engineering control of operator exposure must be used where reasonably practical in addition to the following personal protective equipment: WEAR SUITABLE PROTECTIVE CLOTHING (COVERALLS), SUITABLE PROTECTIVE GLOVES AND FACE PROTECTION (FACESHIELD) when handling the product or contaminated surfaces.

WEAR SUITABLE PROTECTIVE CLOTHING (COVERALLS) during application. However, engineering controls may replace personal protective equipment if a COSHH assessment shows they provide an equal or higher standard of protection.

WHEN USING DO NOT EAT, DRINK OR SMOKE.

WASH HANDS AND EXPOSED SKIN before eating and drinking and after work.

Environmental Protection

Do not contaminate water with the product or its container. Take extreme care to avoid drift onto crops and non-target plants outside the target area. To protect aquatic organisms respect an unsprayed buffer zone to surface water bodies in line with LERAP requirement. DO NOT ALLOW DIRECT SPRAY from horizontal



boom sprayers to fall within 5 m of the top of the bank of a static or flowing waterbody, unless a Local Environmental Risk Assessment for Pesticides (LERAP) permits a narrower buffer zone, or within 1 m of the top of a ditch which is dry at the time of application. Aim spray away from water. This product qualifies for inclusion within the Local Environment Risk Assessment for Pesticides (LERAP) scheme. Before each spraying operation from a horizontal boom sprayer and broadcast air-assisted sprayer, either a LERAP must be carried out in accordance with CRD published guidance or the statutory buffer zone must

be maintained. The results of the LERAP must be recorded and kept available for inspection for three years to any person entitled to exercise enforcement powers under or in connection with the Food and Environment Protection Act (as amended). (An electronic record will satisfy the requirement for a written record, providing it is similarly available for inspection and can be copied).

Storage and Disposal

KEEP IN ORIGINAL CONTAINER, tightly closed in a safe place.

KEEP OUT OF REACH OF CHILDREN.

KEEP AWAY FROM FOOD, DRINK AND ANIMAL FEEDING STUFFS.

WASH OUT CONTAINER THOROUGHLY and dispose of safely.

PROTECT FROM FROST.

DIRECTIONS FOR USE

IMPORTANT: This information is approved as part of the product label. All instructions within this section must be read carefully in order to obtain safe and successful use of this product.

RESTRICTIONS

DO NOT use Clayton Metropolis on crops undersown with grasses, clover or other legumes or any other broad-leaved crop.

Only one "ALS inhibiting" herbicide can be applied to the same crop in sequence or in tank mixture with Clayton Metropolis. Refer to the table under SEQUENCES & TANK MIXTURES for details.

Clayton Metropolis must not be applied to any crop suffering from stress as a result of drought, waterlogging, pest or disease attack, nutrient deficiency, soil compaction or other factors reducing crop growth.

Because some non-target crops are sensitive to Clayton Metropolis, extreme care is required to avoid drift onto plants outside the target area, or onto ponds, waterways or ditches.

Do not apply Clayton Metropolis when rain is imminent.

Do not apply during periods of frosty weather.

Store in a safe dry place designated as an agrochemical store.

WEEDS CONTROLLED

This product contains mesosulfuron-methyl and iodosulfuron-methyl which are ALS inhibitors, also classified by the Herbicide Resistance Action Committee as 'Group B'. Use only as part of a resistance management strategy that includes cultural methods of control and does not use ALS inhibitors as the sole chemical method of grass-weed control. Strains of some annual grasses (e.g. black-grass, wild oats and Italian rye-

grass) have developed resistance to herbicides which may lead to poor control. A strategy for preventing and managing resistance should be adopted. Guidelines have been produced by the Weed Resistance Action Group and copies are available from the HGCA, CPA, your distributor, crop advisor or product manufacturer.

Weeds and Susceptibility

Black-grass (sensitive) - Post-emergence to GS 39

Black-grass (Resistant* - EMR) - Post-emergence to GS 29

Wild oats - Post-emergence to GS 29

Annual meadow-grass - Post-emergence to GS 31

Rough-stalked meadow grass - Post-emergence to GS 31

Perennial rye-grass (from seed) - Post-emergence to GS 31

Italian rye-grass - Post-emergence to GS 30

Common chickweed - Post-emergence to GS 18 (8 expanded true leaves)

Mayweeds - Post-emergence to GS 18 (8 expanded true leaves)

* Situations where Enhanced Metabolism Resistance (EMR) has been confirmed as RR or RRR by a resistance test or where a significant reduction in performance of other herbicides has been noted previously.

Clayton Metropolis is readily translocated within the target weed, inhibiting growth within hours of application. The actual time taken for herbicidal symptoms to appear and death varies between weed species, timing of application and weather conditions. In some cases symptoms may not be apparent for up to 4 weeks. Optimum grass weed control will be obtained when all grass weeds are emerged at spraying. Weeds germinating after application will not be controlled.

Due to the potential for yield loss without prior signs of crop phytotoxicity, avoid use of Clayton Metropolis to control light infestations of grass weeds.

As Clayton Metropolis is active primarily via foliar uptake good spray coverage of the target weed is essential for optimal efficacy. For optimal activity, apply when weather conditions promote active weed growth. Clayton Metropolis controls emerged weeds on all soil types.

Clayton Metropolis has a moderate residual life in soil under normal conditions. As residual activity is important for optimal activity, avoid application under very dry conditions on to very dry soil. Residual efficacy will be enhanced where seedbeds are fine and moist. High soil temperatures and cloddy seedbeds may reduce the residual efficacy of Clayton Metropolis.

The presence of enhanced metabolism herbicide resistant populations of Italian rye-grass may lead to unacceptable levels of control. To reduce the risk of developing resistance or where resistance to sulfonyleurea herbicides is suspected, applications should be made to young, actively growing weeds.

Key aspects of the Clayton Metropolis resistance management strategy are:

ALWAYS follow WRAG guidelines for preventing and managing herbicide resistant grass and broad-leaved weeds.

DO NOT use Clayton Metropolis as a stand-alone treatment for black-grass, rye-grass or common chickweed. Use only in tank mixture or in sequence with herbicides with non-ALS modes of action.

IDEALLY apply Clayton Metropolis as early as possible and before GS 31 of grass weeds.

DO NOT use Clayton Metropolis as the sole means of grass weed or broad-leaved weed control in successive crops.

ALWAYS use grass and broad-leaved weed herbicides with non-ALS modes of action throughout the cropping rotation.

ALWAYS monitor weed control effectiveness and investigate any odd patches of poor grass or broad-leaved weed control. If unexplained contact your agronomist who may consider a resistance test appropriate.

CROP SPECIFIC INFORMATION

Winter Wheat

Apply via a horizontal boom sprayer at a rate of 1.2 L/ha. Apply in 100-300 L/ha as a FINE to MEDIUM spray (BCPC category).

Use application techniques which ensure good weed coverage and crop penetration, using flat fan nozzles. Ensure that spray swaths do not overlap.

Always use Clayton Metropolis in mixture with a suitable authorised adjuvant – eg ADJ: 0617 at a rate of 1.0 L/ha.

Only one application of Clayton Metropolis should be made to the crop.

For use on all varieties of winter wheat.

Apply from the 2 leaf stage (GS 12) of the crop up to flag leaf ligule just visible (GS 39).

SEQUENCES & TANK MIXTURES

Only one "ALS inhibiting" herbicide can be applied to the same crop in sequence or in tank mixture with Clayton Metropolis.

Clayton Metropolis may be applied to the same crop in sequence or in tank mixture with one of a specified number of "ALS inhibiting" herbicides. Check the relevant label for suitability, rates of use or seek advice from your qualified agronomist.

To avoid increased selection of herbicide resistant black-grass, a sequence of Clayton Metropolis may be used with a suitably approved product containing flupyr-sulfuron-methyl if it has been applied in tank-mix with a suitable residual partner e.g. pendimethalin and before black-grass emergence.

These sequences and tank mixtures must only be applied in accordance with label recommendations for every product in the sequence or tank mixture and only if used in conjunction with a robust non-ALS inhibitor autumn herbicide programme. This is particularly important where both components are applied in the spring. Aside from those mentioned above, do not use in sequence or tank mixture with any other ALS inhibitors, such as sulfonylureas, unless specifically permitted on the label of those products.

As part of the Herbicide Resistance Management Strategy for Clayton Metropolis, do not apply herbicides MAPP16495 or MAPP16494 in sequence with Clayton Metropolis for the control of common chickweed in order to avoid increased selection for herbicide resistant individuals.

FOLLOWING CROPS and CROP FAILURE

Only winter wheat, winter barley or winter oilseed rape may be sown in the year of harvest to succeed a winter wheat crop treated with Clayton Metropolis. Spring wheat, spring barley, spring oilseed rape and sugar beet may be drilled in the spring following harvest of the Clayton Metropolis treated winter wheat crop. Plough prior to planting crops of oilseed rape, otherwise crop damage may occur. In the event of crop failure for any reason, sow only winter wheat in the same cropping season as an application of Clayton Metropolis. Where Clayton Metropolis is applied in sequence or tank mixture with other permitted "ALS inhibiting" herbicides, always follow the most restrictive label with regard to following crops.

MIXING

Add the recommended quantity of Clayton Metropolis to the spray tank half-filled with the required quantity of clean water.

Add the remainder of the water with the sprayer agitation system in operation.

Maintain agitation during mixing and loading and until spraying is complete.

Do not leave the sprayer standing with chemical in it.

To avoid subsequent damage to crops other than winter wheat it is important that the spray tank, boom, hoses, filters and nozzles are thoroughly washed out to remove all traces of Clayton Metropolis immediately after spraying using a suitable proprietary sprayer cleaner according to the label instructions for that product.

=====