


CLAYTON ABBA

MAPP 18559

Contains 18 g/l (1.84 % w/w) abamectin with cyclohexanol in an emulsifiable concentrate

For the control of the two-spotted red spider mite in tomatoes, cucumbers and in ornamental plant production; additionally Western Flower Thrips in cucumbers and ornamentals and leaf miner in tomatoes.

	<p>Clayton Abba contains abamectin with cyclohexanol Signal Word :Warning</p> <p>Hazard Statements : Harmful if swallowed Causes serious eye irritation May cause damage to the nervous system through prolonged or repeated exposure</p> <p>DANGEROUS FOR THE ENVIRONMENT Very toxic to aquatic organisms, may cause long-term adverse effects in the aquatic environment</p>
---	--

The Control of Substances Hazardous to Health (COSHH) Regulations may apply to the use of this product at work.

IMPORTANT INFORMATION

FOR USE ONLY AS A <u>HORTICULTURAL INSECTICIDE/ACARICIDE</u>			
Crop	Maximum individual dose of product	Maximum number of treatments	Latest time of application
Tomato (protected)	See Other Specific	2 pre-bloom or fruit set and	3 days before harvest
Cucumber (protected)	Restriction 1	4 post-bloom or fruit set	
Ornamental plant	See Other Specific	- production	Restriction 1

Other specific restrictions

1. The maximum concentration must not exceed 50 ml product per 100 litres water.
2. Protected tomatoes and cucumbers which are in flower or have set fruit must not be treated between 1 November and end-February.
3. Cherry tomatoes must not be treated.
4. KEEP UNPROTECTED PERSONS OUT OF TREATED AREAS until the spray has dried.

READ THE LABEL BEFORE USE. USING THIS PRODUCT IN A MANNER THAT IS INCONSISTENT WITH THE LABEL MAY BE AN OFFENCE. FOLLOW THE CODE OF PRACTICE FOR USING PLANT PROTECTION PRODUCTS

SAFETY PRECAUTIONS Operator protection

Engineering control of operator exposure must be used where reasonably practicable in addition to the following personal protective equipment:

WEAR SUITABLE PROTECTIVE CLOTHING (COVERALLS, APRON), SUITABLE PROTECTIVE GLOVES, RUBBER BOOTS, FACE PROTECTION (FACESHIELD) AND SUITABLE RESPIRATORY PROTECTIVE EQUIPMENT* when handling the concentrate and when spraying.

* Disposable filtering facepiece respirator to at least EN149 FFP2 or equivalent.

WEAR SUITABLE PROTECTIVE CLOTHING (COVERALLS), SUITABLE PROTECTIVE GLOVES, RUBBER BOOTS AND FACE PROTECTION (FACESHIELD) when handling contaminated surfaces or freshly-treated material.

However, engineering controls may replace personal protective equipment if a COSHH assessment shows they provide an equal or higher standard of protection. Do not breathe dust/fume/gas/mist/vapours/spray WHEN USING DO NOT EAT, DRINK OR SMOKE.

WASH HANDS AND EXPOSED SKIN before eating, drinking or smoking and after work.

WASH ALL PROTECTIVE CLOTHING thoroughly after use, especially the insides of gloves.

IN CASE OF ACCIDENT OR IF YOU FEEL UNWELL, seek medical advice immediately (show the label where possible). IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing. : Get medical advice/attention

Environmental protection

Do not contaminate water with the product or its container. Do not clean application equipment near surface water. Avoid contamination via drains from farmyards and roads. Use appropriate containment to avoid environmental contamination.

HIGH RISK TO BEES. DO NOT apply to crops in flower or to those in which bees are actively foraging. Do not apply when flowering weeds are present.

Storage and disposal

KEEP AWAY FROM FOOD, DRINK AND ANIMAL FEEDING STUFFS.

KEEP IN ORIGINAL CONTAINER, tightly closed, in a safe place under lock and key.

KEEP OUT OF REACH OF CHILDREN.

This material and its container must be disposed of in a safe way.

DO NOT RE-USE CONTAINER for any purpose.

RINSE CONTAINER THOROUGHLY by using an integrated pressure rinsing device or manually rinsing three times. Add washings to sprayer at time of filling and dispose of the container safely.

**Clayton Plant Protection (UK) Ltd.,
Unit F10, Bracetown Business Park
CLONEE, Dublin 15 Ireland.**

Contents: **1 litre e**

PROTECT FROM FROST

Tel: (00 353) 1 8210127

Batch No:

Conditions of Supply: all goods supplied by us are of high quality and we believe them to be correct but, as we cannot exercise control over their storage, handling, mixing or use, or weather conditions before, during and after application which may affect the performance of the goods, all conditions and warranties, statutory or otherwise, as to the quality or fitness for any purpose of our goods are excluded, and no responsibility will be accepted by us or resellers for any failure in performance, damage or injury whatsoever arising from their storage, handling, application or use. These conditions cannot be varied by our staff or agents whether or not they supervise or assist in the use of such goods.

DIRECTIONS FOR USE

IMPORTANT: This information is approved as part of the Product Label. All instructions within this section must be read carefully in order to obtain safe and successful use of this product.

ACTIVITY

Clayton Abba, containing abamectin, is an insecticide and acaricide with contact and stomach activity. It is not systemic but exhibits translaminar movement. Clayton Abba is approved for the control of the two-spotted red spider mite in tomatoes, cucumbers and in ornamental plant production; in addition, when used at the higher rate, nymphs of the Western Flower Thrips are controlled in cucumbers and ornamentals as well as leaf miner larvae in tomatoes. Susceptible pests become inactive soon after treatment but their death may not occur until several days have elapsed following treatment.

PROTECTED TOMATOES AND CUCUMBERS

Crop	Pest	Dose	Treatment advice
Protected tomatoes excluding cherry tomatoes.	Two-spotted spider mite, <i>Tetranychus urticae</i> .	25 ml per 100 litres water.	Treat as soon as pest is seen before damage occurs. Repeat after 7 days if necessary.
	Leaf miner larvae, <i>Liriomyza bryoniae</i> .	50 ml per 100 litres water.	Treat as soon as pest is seen or early leaf mining has occurred. Repeat after 7 days if necessary.
Protected cucumbers	Two-spotted spider mite, <i>Tetranychus urticae</i> .	25 ml per 100 litres water.	Treat as soon as pest is seen before damage occurs. Repeat after 7 days if necessary.
	Nymphs of Western flower thrip, <i>Frankliniella occidentalis</i>	50 ml per 100 litres water.	Treat as soon as pest is seen. Repeat after 7 days and again after 14 days if necessary and permissible.

Restrictions

- 1) Crops in flower or with set fruit may only be treated from 1st March to 31st October inclusive. Do not treat flowering crops or crops in fruit between 1st November and 29th February.
- 2) 2 applications pre-bloom or fruit set and 4 post-bloom or fruit set may be made per crop in total.

ORNAMENTAL PLANT PRODUCTION

Crop	Pest	Dose	Treatment advice
Protected and outdoor ornamentals and flower crops.	Two-spotted spider mite, <i>Tetranychus urticae</i> .	25 ml per 100 litres water.	Treat as soon as pest is seen before damage occurs. Repeat after 7 days if necessary.
	Nymphs of Western flower thrip, <i>Frankliniella occidentalis</i>	50 ml per 100 litres water.	Treat as soon as pest is seen. Repeat after 7 days and again after 14 days if necessary.

Restrictions

- 1) Do not apply to crops in flower.
- 2) Verify the safety of the treatment on all new or unusual species and varieties to which one is unaccustomed before proceeding to treat large numbers of plants.
- 3) **Do not treat** ferns or Shasta daisies.

Caution

Some leaf spotting or staining may occur on some plant species. Leaf marking has occurred on Carnation, Kalanchoe and Begonia.

BENEFICIAL INSECTS

Clayton Abba is not recommended for use in integrated pest management (IPM) systems.

When bumble bees are employed as pollinators in tomato crops, it is recommended that their introduction is delayed for at least 24 hours after treatment of the crop.

REPEATED USE

Too frequent use of Clayton Abba is not recommended. After two consecutive applications have been made, it advisable to introduce an alternative product with a different mode of action into spray programmes in order to minimise the selection pressure for resistant strains.

APPLICATION

Apply as a high-volume spray through a hand lance fitted with appropriate nozzles using a conventional high-volume hydraulic, powered or knapsack, sprayer. Thoroughly cover the stems and both sides of the crop leaves with spray to just before the point of run-off.

MIXING

Half-fill the spray tank with clean water and put under agitation. Shake the product container thoroughly before pouring and mix in the required volume of Clayton Abba through the top filter or filling device. Fill the spray tank to the required level with more clean water and allow thorough mixing under agitation before use. Spray immediately after mixing and keep under agitation until sprayed out. Do not allow mixed spray to stand in the tank for long periods.

COMPATIBILITY

The addition of any adjuvant, sticker or wetting agent is neither required nor recommended; do not mix with any adjuvant.

EQUIPMENT MAINTENANCE

Immediately after use wash the spraying machine and all utensils thoroughly with clean water and a wetting agent recommended for the cleaning of application equipment.