

# AZOFIN PLUS

A suspension concentrate formulation containing 250 g/L (23.2% w/w) of azoxystrobin. A broad-spectrum fungicide to be used on asparagus (outdoor), barley, beans (broad beans, fresh green beans, field beans), bulb onions, carrots, endive, escarole, frisee, garlic, leeks, lettuce, lupins, oats, oilseed rape (winter and spring), outdoor *Brassicaceae* (broccoli, calabrese, Brussels sprout, cabbage, cauliflower, winter greens kale, spring greens collard), peas (combining, vining and fresh varieties), potatoes (in furrow treatment and foliar spray), radicchio chicory, rye, shallots, strawberries, triticale, wheat.

MAPP 18552



**AZOFIN PLUS. A suspension concentrate formulation containing 250 g/L of azoxystrobin. Contains 1,2-benzisothiazol-3-one. May produce an allergic reaction.**

**WARNING :** Harmful if inhaled.  
Very toxic to aquatic life with long lasting effects.



Avoid breathing dust/fumes/gas/mist/vapours/spray. Use only outdoors or in a well-ventilated area. **IF INHALED:** remove person to fresh air and keep comfortable for breathing. Call a **POISON CENTRE/doctor** if you feel unwell. Collect spillage. Dispose of contents/container to a licensed hazardous-waste disposal contractor or collection site except for empty clean containers which can be disposed of as non-hazardous waste.

To avoid risks to human health and the environment, comply with the instructions for use.

IMPORTANT INFORMATION : FOR USE ONLY AS AN AGRICULTURAL/HORTICULTURAL FUNGICIDE				
Crops	Max individual dose (L product/ha)	Max number of treatments (per crop)	Minimum spray interval (days)	Latest time of application
Asparagus (outdoor)	1.0	2	10	Before senescence
Barley, Oats	1.0	2	14	Before 61 (flowering beginning)
Beans, broad (fresh, edible podded pea, vining pea)	1.0	2	14	14 days before harvest
Dwarf French bean	1.0	2	14	7 days before harvest
Beans- field, lupins	1.0	2	21	35 days before harvest
**Broccoli/calabrese, Brussels sprout, cabbage, cauliflower, kale (winter greens), collard (spring greens) – all outdoor	1.0	2	12	14 days before harvest
Bulb onion, garlic, shallots	1.0	3	7	14 days before harvest
Carrot	1.0	3	7	14 days before harvest

Leek	1.0	3	12	21 days before harvest
**Lettuce, endive, (outdoor and protected)	1.0	2	7	14 days before harvest
Oilseed rape (winter and spring)	1.0	2	21	21 days before harvest
Peas, combining	1.0	2	14	35 days before harvest
Potato, in furrow	3.0	1	-	At planting
Potato, foliar spray	0.5	3	7	7 days before harvest
Strawberry (outdoor and protected)	1.0	3	7	3 days before harvest
Wheat, rye and triticale	1.0	2	14	Before GS71 (before watery ripe stage)

**Other specific restrictions:** To reduce the risk of resistance developing in target diseases, the number of applications of product containing QoI fungicides made to cereal crops must not exceed two. When used in a protected situation other than 'permanent protection with full enclosure', aquatic buffer zones in line with LERAP requirements must be observed. \*\* A maximum total dose of 500g azoxystrobin must not be exceeded within a 12 month period on the same field.

**READ THE LABEL BEFORE USE. FOLLOW THE CODE OF PRACTICE FOR USING PLANT PROTECTION PRODUCTS. USING THIS PRODUCT IN A MANNER THAT IS INCONSISTENT WITH THE LABEL MAY BE AN OFFENCE.**

**Approval Holder**

Finchimica S.p.A, via Lazio, 13, 25025 Manerbio (BS), Italy  
**Marketing company in UK**  
**Clayton Plant Protection (UK) Ltd. Unit F10,**  
**Bracetown Business Park**  
**Clonee, Co. Meath. Ireland Tel:**  
(00 353) 1 8210127  
[www.claytonpp.com](http://www.claytonpp.com) Email: [info@claytonpp.com](mailto:info@claytonpp.com)

**STORE IN A COOL DRY PLACE**

**PROTECT FROM FROST**

**SHAKE WELL BEFORE USE**

**Contents : xx L e**

**UN3082**

Batch no:

The (COSHH) Control of Substances Hazardous to Health Regulations may apply to the use of this product at work.

**SAFETY PRECAUTIONS Operator protection**

Wash splashes from skin or eyes immediately.

Wash hands and exposed skin before meals and after work.

Do not breathe spray.

**Environmental protection**

Avoid drift onto non-target plants.

Do not contaminate water with this product or its container.

Do not clean application equipment near surface water. Avoid contamination via drains from farmyards and roads.

To protect aquatic life, for uses on crops broccoli, calabrese, Brussels sprouts, cabbage, cauliflower, collards, lettuce and kale, the maximum total dose applied must not exceed 500 g azoxystrobin per hectare per year. To protect aquatic organisms, respect an unsprayed buffer zone to surface water bodies in line with LERAP requirements.



DO NOT ALLOW DIRECT SPRAY from horizontal boom sprayers to fall within 5 m of the top of the bank of a static or flowing water body, unless a Local Environment Risk Assessment for Pesticides (LERAP) permits a narrower buffer zone, or within 1 m of the top of a ditch which is dry at the time of application.

DO NOT ALLOW DIRECT SPRAY from hand held sprayers to fall within 1 m of the top of the bank of a static or flowing waterbody. Aim spray away from water.

AZOFIN PLUS qualifies for inclusion within the LERAP (Local Environmental Risk Assessment for Pesticides) Scheme. Before each spraying operation from a horizontal boom sprayer, either a LERAP must be carried out (according to CRD published guidance) or the statutory buffer zone must be maintained. The results must be recorded and kept available for inspection for 3 years.

**Storage and disposal**

KEEP IN ORIGINAL CONTAINER tightly closed, in a safe place.

WASH OUT CONTAINER THOROUGHLY, empty washings into spray tank and dispose of safely. DISPOSE OF THE PRODUCT AND ITS CONTAINER in a safe way.

**DIRECTIONS FOR USE**

**IMPORTANT:** This information is approved as part of the Product Label. All instructions of this section must be carefully observed in order to obtain safe and successful use of this product.

## GENERAL INFORMATION

AZOFIN PLUS contains azoxystrobin, a broad-spectrum fungicide from the strobilurin group that inhibits fungal respiration. It should always be used in mixture with fungicides with other modes of action.

It has systemic, translaminar and protectant properties. AZOFIN PLUS is best used as a protective treatment or during early stages of disease establishment. In cereals, the length of disease control is generally about four to six weeks during the period of active stem elongation, but can be more when applied at flag leaf/ear emergence. AZOFIN PLUS is approved for application to asparagus (outdoor), barley, beans (broad beans, fresh green beans, field beans), bulb onions, carrots, endive, escarole, frisee, garlic, leeks, lettuce, lupin, oats, oilseed rape (winter and spring), outdoor *Brassicaceae* (broccoli, calabrese, Brussels sprout, cabbage, cauliflower, winter greens kale, spring greens collard), peas (combining), potatoes (in furrow treatment and foliar spray), radicchio chicory, rye, shallots, strawberries, triticale, wheat.

## RESTRICTIONS

Certain apple varieties are highly sensitive to AZOFIN PLUS. As a precaution AZOFIN PLUS should not be applied when there is a risk of spray drift onto neighbouring apple crops. Spray equipment used to apply AZOFIN PLUS to other crops should not be used to treat apples. Apply AZOFIN PLUS under good growing conditions with adequate soil moisture. Avoid poor growing conditions which may give less reliable results.

CONSULT PROCESSORS BEFORE USING ON CROPS GROWN FOR PROCESSING.

DO NOT USE on soils with high organic matter content as control may be reduced.

## RESISTANCE

AZOFIN PLUS is an azoxystrobin-based formulation. Azoxystrobin is a member of the QoI cross resistance group. To obtain the best results, AZOFIN PLUS should be used preventatively.

To reduce the risk of resistance developing in target diseases, the total number of applications of product containing QoI fungicides made to any cereal crop must not exceed two.

Use AZOFIN PLUS as part of an Integrated Crop Management (ICM) strategy, including fungicides with a different mode of action.

AZOFIN PLUS must be applied with due regard to current FRAG-UK or FRAC guidelines for the relevant crop for QoI compounds.

## DISEASES CONTROLLED

Barley	Brown Rust ( <i>Puccinia hordei</i> ). Leaf Blotch ( <i>Rhynchosporium secalis</i> ) - reduction Net Blotch ( <i>Pyrenophora teres</i> ). Can reduce severity of Take-all ( <i>Gaeumannomyces graminis var. Tritici</i> )
Oats	Crown Rust ( <i>Puccinia coronata</i> )
Rye and Triticale	Brown Rust ( <i>Puccinia recondita</i> ). Leaf Blotch ( <i>Rhynchosporium secalis</i> ) - reduction Can reduce severity of Take-all ( <i>Gaeumannomyces graminis var. Tritici</i> )
Wheat	Brown Rust ( <i>Puccinia recondita</i> ). Ear Diseases ( <i>Cladosporium</i> , <i>Alternaria</i> ). Glume Blotch ( <i>Leptosphaeria (syn. Septoria) nodorum</i> ). Yellow Rust ( <i>Puccinia striiformis</i> ) Can reduce severity of Take-all ( <i>Gaeumannomyces graminis var. Tritici</i> )
Oilseed rape	Dark Leaf and Pod Spot ( <i>Alternaria spp.</i> ). Sclerotinia stem rot ( <i>S. sclerotiorum</i> ) – moderate control
Peas	Downy mildew ( <i>Peronospora viciae</i> ) - reduction Leaf and Pod Spot ( <i>Ascochyta pisi</i> ) – useful control. If used against leaf and pod spot, some control of Grey mould ( <i>Botrytis cinerea</i> ) and <i>Mycosphaerella</i> blight may be achieved.
Bulb Onions, Garlic and Shallot	Downy mildew ( <i>Peronospora destructor</i> ) – moderate control
Leeks	Leaf rust ( <i>Puccinia porri</i> ). Purple blotch ( <i>Alternaria porri</i> ) – moderate control White tip ( <i>Phytophthora porri</i> ) – moderate control
Carrots	<i>Alternaria</i> leaf blight ( <i>Alternaria dauci</i> ) Powdery mildew ( <i>Erysiphe polygoni</i> )
Asparagus	Stemphylium ( <i>Stemphylium botryosum</i> ) – moderate control. Rust ( <i>Puccinia asparagi</i> ) – moderate control
Broad Beans, Field Beans	Rust ( <i>Uromyces fabae</i> )
Brussels Sprouts, Cabbage, Cauliflower, Kale (Winter Greens), Collards (Spring Greens), Broccoli and Calabrese	White blister ( <i>Albugo candida</i> ) – moderate control Ring spot ( <i>Mycosphaerella brassicicola</i> ) – moderate control <i>Alternaria (Alternaria brassicae and Alternaria brassicicola)</i> – moderate control
Lupins	Rust ( <i>Uromyces spp.</i> ) – Qualified Use recommendation
Potatoes	Stem canker and Black scurf ( <i>Rhizoctonia solani</i> ) – reduction in furrow only Black dot ( <i>Colletotrichum coccodes</i> ) – reduction in furrow use only Early blight ( <i>Alternaria solani</i> ) - moderate control foliar use only

Strawberry	Powdery mildew ( <i>Podosphaera macularis</i> ) – moderate control Anthracnose ( <i>Colletotrichum acutatum</i> ) - Qualified Use recommendation
Lettuce, Endive	Downy mildew ( <i>Bremia spp.</i> )

### **CROP SPECIFIC INFORMATION Barley, oats, wheat, rye, triticale**

Apply AZOFIN PLUS (max 2 applications per crop with a minimum spray interval of 14 days) at a rate of 1.0 L/ha, in a minimum water volume of 200 L/ha. Apply at earliest stage of disease development or as protectant treatment following a disease risk assessment or the use of appropriate decision support systems.

Winter and spring wheat, rye and triticale can be treated from BBCH 30-69.

Winter and spring barley and winter and spring oats can be treated from BBCH 30-59.

For protection against ear disease (*Cladosporium* and *Alternaria*) apply AZOFIN PLUS at ear emergence.

When used to control the listed foliar diseases, AZOFIN PLUS applied at the first or second node stage of the crop can reduce the severity of Take-all infection.

When used on cereals crops, AZOFIN PLUS must be mixed with a product containing a fungicide from a different cross resistance group that is recommended for same target disease control.

AZOFIN PLUS must be used as part of an Integrated Crop Management (ICM) strategy incorporating other methods of control.

Do not apply more than 2 foliar applications of QoI-containing products to any cereal crop. Failure to follow resistance management action may result in reduced levels of disease control. Disease control may be reduced if strains of other pathogens less sensitive to azoxystrobin develop. For details, please refer to FRAG-UK guidelines for QoI-compounds.

### **Peas (combining and fresh)**

If crops are to be processed then processors should be consulted before treating with AZOFIN PLUS. Apply AZOFIN PLUS (max 2 applications per crop with a minimum spray interval of 14 days), at a rate of 1.0 L/ha, in a minimum water volume of 200 litres per hectare.

Before the application, ensure the crop is free from any stress caused by agronomic and/or environment effects.

If needed, wax level in peas can be checked using the Crystal Violet test.

Apply at earliest stage of disease development (recommended timings are BBCH 17-72).

A second treatment may be required if disease pressure remains high – especially in combining peas.

AZOFIN PLUS should always be used at the first sign of disease infection or when a predictive assessment shows conditions favourable for disease development. For optimum disease control apply AZOFIN PLUS before infection or as soon as disease is first seen in the crop. Always inspect crops to assess disease development immediately before spraying. Best results will be achieved from applications made in the earliest stage of disease development or as a protectant treatment following a disease risk assessment or the use of appropriate decision support systems.

AZOFIN PLUS must be applied with due regard to current FRAG-UK guidelines for QoI compounds. Do not exceed 2 applications to peas.

### **Bulb onions, leeks, carrots**

Apply AZOFIN PLUS (max 3 applications per crop, with a minimum spray interval of 7 days for bulb onion, carrots, shallot and garlic and 12 days for leeks) at a rate of 1.0 L/ha, in a minimum water volume of 200 litres per hectare.

Always inspect crops to assess disease development immediately before spraying. Best results will be achieved from applications made in the earliest stage of disease development or as a protectant treatment following a disease risk assessment or the use of appropriate decision support systems.

#### **Recommended application timings are:**

Bulb onions, garlic and shallot, BBCH 14-48. Leeks BBCH 16-48. Carrots BBCH 16-49.

Latest time of application is 14 days before harvest for bulb onion, carrots, shallot and garlic, and 21 days for leeks.

The product must be used as part of an Integrated Crop Management (ICM) strategy incorporating other methods of control.

AZOFIN PLUS must be applied with due regard to current FRAC guidelines for QoI compounds.

Do not use more than 3 applications of AZOFIN PLUS per crop.

For optimum downy mildew control in bulb onions, garlic and shallot a 7 to 10 day spray interval should be maintained. Applications to established downy mildew infection are unlikely to give reliable control.

To avoid the likelihood of resistance developing, applications of AZOFIN PLUS should be made with due regard to current FRAC guidelines for QoI compounds as illustrated below in the following table:

Total number of fungicide spray applications per crop	1	2	3	4	5	6	7	8	9	10	11	≥ 12
Maximum recommended solo QoI fungicide sprays	1	1	2	2	2	2	2	3	3	3	3	4
Maximum recommended QoI fungicide sprays in mixture	1	2	2	2	2	3	3	4	4	4	4	4

No more than 3 applications of AZOFIN PLUS are permitted per crop. Refer to the FRAC website for updates on recommendations for resistance management.

If crops are to be processed then processors should be consulted before treating with AZOFIN PLUS.

### Asparagus (outdoor)

Apply AZOFIN PLUS (max 2 applications per crop, with a minimum spray interval of 10 days) at a rate of 1.0 L/ha.

Recommended water volumes:

- minimum 600 litres per hectare for application through tractor mounted spraying equipment -
- minimum 200 litres per hectare for hand-held spraying equipment.

Before the application, ensure the crop is free from any stress caused by agronomic and/or environment effects. Apply at earliest stage of disease development or as protectant treatment following a disease risk assessment.

Recommended treatment is BBCH 41-89, note the earliest time of application of AZOFIN PLUS is after commercial cutting after the harvest season. If a new bed is established then treatment should not take place within 3 weeks of transplanting out the crowns.

The latest application timing is the end of September or before crop senescence, whichever is earliest.

AZOFIN PLUS contains azoxystrobin a member of the QoI cross resistance group. AZOFIN PLUS should be used preventatively and should not be relied on for its curative potential. Disease control may be reduced if strains of pathogens less sensitive to azoxystrobin develop.

To avoid the likelihood of resistance developing, applications of AZOFIN PLUS should be made with due regard to current FRAC guidelines for QoI compounds as illustrated below in the following table:

Total number of fungicide spray applications per crop	1	2	3	4	5	6	7	≥8
Maximum recommended solo QoI fungicide sprays	1	1	2	2	2	2	2	3
Maximum recommended QoI fungicide sprays in mixture	1	2	2	2	2	3	3	3

No more than 2 applications of AZOFIN PLUS are permitted per crop. Refer to the FRAC website for updates on recommendations for resistance management.

### Beans and Lupins

Apply AZOFIN PLUS (max 2 applications per crop, with the second treatment required in cases of high disease pressure at a minimum interval of 21 days for field bean and lupin, and 14 days for broad bean and fresh bean) at a rate of 1.0 L/ha, in a minimum water volume of 150 litres per hectare. Apply at earliest stage of disease development (recommended BBCH 60-69 for field beans, BBCH 17-72 for fresh beans and lupins) or as protectant treatment following a disease risk assessment.

#### Latest time of application:

Broad beans: 14 days before harvest. Fresh beans: 7 days before harvest. Field beans, lupins: 35 days before harvest. Before applying AZOFIN PLUS, ensure the crop is free from any stress caused by environmental or agronomic effects. Always inspect crops to assess disease development immediately before spraying. Best results will be achieved from applications made in the earliest stage of disease development or as a protectant treatment following a disease risk assessment or the use of appropriate decision support systems".

To avoid the likelihood of resistance developing, application of AZOFIN PLUS should be made with due regard to current FRAG-UK guidelines for QoI compounds. Do not make more than two applications of AZOFIN PLUS. Use AZOFIN PLUS as part of an Integrated Crop Management (ICM) strategy incorporating other methods of control, including where appropriate other fungicides with a different mode of action.

### Broccoli, Calabrese, Brussels sprouts, Cabbage, Cauliflower, Kale (Winter Greens), Collards (Spring Greens)

Apply AZOFIN PLUS (max 2 applications per brassica crop, with the second treatment to be used just in case of high disease pressure, with an interval of 12 days between applications) at a rate of 1.0 L/ha, in a minimum water volume of 250 litres per hectare.

Always inspect crops to assess disease development immediately before spraying.

Before the application, ensure the crop is free from any stress caused by agronomic and/or environment effects. Apply at earliest stage of disease development or as protectant treatment (recommended timing BBCH 16-49).

Latest time of application is 14 days before harvest.

A maximum total of 500 g of azoxystrobin must not be exceeded within 1 year period on the same field.

AZOFIN PLUS must be applied with due regard to current FRAG guidelines for QoI compounds. Do not use more than 2 applications of AZOFIN PLUS per crop.

## Potatoes

Use AZOFIN PLUS only on potato crops adhering to good rotation practices.

If crop is to be processed then processors should be consulted before treating with AZOFIN PLUS.

**Foliar application against early blight:** Before applying AZOFIN PLUS, ensure the crop is free from any stress caused by environmental or agronomic effects. Always inspect crops to assess disease development immediately before spraying. Best results will be achieved from applications made in the earliest stage of disease development or as a protectant treatment following a disease risk assessment or the use of appropriate decision support systems. Apply AZOFIN PLUS (max 3 applications with a minimum spray interval of 7 days) at a rate of 0.5 L/ha, in a minimum water volume of 200 litres per hectare. Where a crop of potatoes is destined for processing, consult processors before treating with AZOFIN PLUS.

Crop may be treated from BBCH 51-85 – latest application timing is 7 days before harvest.

The risk of resistance developing to AZOFIN PLUS in *Alternaria solani* is considered to be moderate. To avoid the likelihood of resistance developing, application of AZOFIN PLUS should be made with due regard to current FRAG-UK guidelines for QoI compounds. Use AZOFIN PLUS as part of an Integrated Crop Management (ICM) strategy incorporating other methods of control, including where appropriate other fungicides with a different mode of action.

**In furrow application against stem canker, black scurf and black dot:** Apply AZOFIN PLUS (max 1 application) at a rate of 3.0 L/ha in a water volume of 50-150 litres per hectare.

AZOFIN PLUS must be applied as an in-furrow application made at the time of planting. Where AZOFIN PLUS is applied as an in-furrow application, it is important to direct the spray into the planting furrow and not onto the seed tuber.

Application should ensure that the AZOFIN PLUS is applied to soil around the tuber.

With in-furrow application, always target the soil and not the seed tuber in order to minimise any possible delay in emergence.

Wherever possible, use properly chitted seed or cold-stored seed which has not started to sprout. Using seed which has just broken dormancy may well result in emergence delays.

Using AZOFIN PLUS following earlier applications of imazalil, pencycuron or imazalil/pencycuron is likely to lead to a check in the speed of crop emergence. Effects are usually, but not always, outgrown.

Do not use AZOFIN PLUS on high organic matter soils as the product will not be effective.

Where a crop of potatoes is destined for processing, consult processors before treating with AZOFIN PLUS.

The risk of resistance developing to AZOFIN PLUS in *Rhizoctonia solani* (Black scurf and Stem canker) and *Colletotrichum coccodes* (Black dot) is considered to be very low. AZOFIN PLUS should only be used in potato crops, which adhere to good rotation practices. To avoid the likelihood of resistance developing to QoI compounds used to control potato late blight, application of AZOFIN PLUS should be made with due regard to current FRAG-UK guidelines for QoI compounds. If an application of AZOFIN PLUS is made, no more than two further QoI treatments should be applied sequentially as the first sprays against late blight before using an alternative product

## Oilseed rape

Before applying AZOFIN PLUS, ensure the crop is free from any stress caused by environmental or agronomic effects. A second treatment may be required if disease pressure remains high.

Apply AZOFIN PLUS (max 2 applications per crop with a minimum spray interval of 21 days) at a rate of 1.0 L/ha, in a minimum water volume of 200 litres per hectare. Apply at earliest stage of disease development or as protectant treatment following a disease risk assessment, the latest time of application is 21 days before harvest.

AZOFIN PLUS must be applied with due regard to current FRAG guidelines for QoI compounds. Do not apply more than 2 applications of AZOFIN PLUS per crop. The product should be used as part of an Integrated Crop Management (ICM) strategy incorporating other methods of control, including where appropriate other fungicides with a different mode of action.

### **Recommendations:**

Use AZOFIN PLUS against *Sclerotinia* as a protectant during flowering (best results from an application timing GS60-GS65, from early to mid-flowering).

Use AZOFIN PLUS against *Alternaria* as a protectant at early pod formation (before the first spots appear on the pods).

## Lettuce, Endive, Chicory (outdoor and protected)

Apply AZOFIN PLUS (max 2 applications per crop, with a minimum spray interval of 7 days between applications for both indoor and outdoor crops) at a rate of 1.0 L/ha, in a minimum water volume of 300 litres per hectare (both outdoor and protected). Before the application, ensure the crop is free from any stress caused by agronomic and/or environment effects. Apply at earliest stage of disease development or as protectant treatment following a disease risk assessment, recommended timing is from BBCH 14-49, the latest time of application is 14 days before harvest.

AZOFIN PLUS must be applied with due regard to current FRAG guidelines for QoI compounds. Do not use more than 2 applications of AZOFIN PLUS per crop. The product should be used as part of an Integrated Crop Management (ICM) strategy incorporating other methods of control, including where appropriate other fungicides with a different mode of action.

A maximum total of 500 g of azoxystrobin must not be exceeded within 1 year period on the same field.

### Strawberry (outdoor and protected)

Apply AZOFIN PLUS (max 3 applications per crop, with a minimum spray interval of 7 days between applications for both indoor and outdoor crops) at a rate of 1.0 L/ha.

Recommended water volumes: - at least 300 litres of water per hectare for outdoor crop;  
- at least 100 litres of water per hectare for indoor crop.

Before the application, ensure the crop is free from any stress caused by agronomic and/or environment effects.

The recommended treatment timing is from BBCH 51-89, the latest time of application is 3 days before harvest. If crops are to be processed, then processors should be consulted before treating with AZOFIN PLUS.

For optimum results apply AZOFIN PLUS as a protectant spray at the beginning of flowering. Two further applications can be made if disease pressure remains high. Application should be made in sequence with other products as part of a fungicide programme during flowering at a minimum interval of 7 days.

Use AZOFIN PLUS as part of an Integrated Crop Management (ICM) strategy incorporating other methods of control, including where appropriate other fungicides with a different mode of action.

To avoid the likelihood of resistance developing, applications of AZOFIN PLUS should be made with due regard to current FRAC guidelines for QoI compounds as illustrated below in the following table:

Total number of fungicide spray applications per crop	1	2	3	4	5	6	7	≥8
Maximum recommended solo QoI fungicide sprays	1	1	2	2	2	2	2	3
Maximum recommended QoI fungicide sprays in mixture	1	2	2	2	2	3	3	3

No more than 3 applications of AZOFIN PLUS are permitted per crop.

### QUALIFIED USE RECOMMENDATION Lupins – Strawberries

The below listed uses are supported by a limited amount of effectiveness data which indicate that the use of AZOFIN PLUS at 1.0 L/ha may provide some useful activity against Rust (*Uromyces spp*) on Lupins and Anthracnose (*Collectotrichum acutatum*) on strawberries. For lupins apply in at least 200L/ha of water, for strawberries 100L/ha for indoor uses and 300L/ha for outdoor uses.

### MIXING AND SPRAYING Application Equipment: conventional crop spraying equipment

**Sprayer Preparation:** Ensure that the sprayer is clean and correctly calibrated. Apply at the water volumes appropriate for the density of the crop, using a medium quality spray (as defined by BCPC guidelines) to guarantee a good coverage.

**Spray Pressure:** at least 2 bar.

**Mixing:** Before mixing AZOFIN PLUS, half-fill the spray tank with clean water and start agitation. Shake the container before opening. Open the AZOFIN PLUS container and add the recommended amount of the product to the sprayer using a filling device, or otherwise by direct adding the product to the spray tank.

Flush thoroughly the empty container and add washings to the sprayer. An integrated pressure rinsing device is suggested; otherwise manually rinse the container three times. Add the remainder of the water and continue agitation during spraying.

Do not leave the spray liquid in the sprayer for prolonged periods. Therefore, never prepare more spray than required.

**Tank cleaning:** After use, wash measuring and spray equipment thoroughly.

**Volume of Water :** Please refer to Crop Specific Information on this booklet. To ensure good coverage, water volumes may need to be increased in dense crops.

### AFTER SPRAYING

Thoroughly wash out sprayer according to manufacturer's guidelines and dispose of washing and clean containers according to DEFRA Code of Practice and local water authority guidelines.

**Conditions of Supply:** all goods supplied by us are of high quality and we believe them to be correct but, as we cannot exercise control over their storage, handling, mixing or use, or weather conditions before, during and after application which may affect the performance of the goods, all conditions and warranties, statutory or otherwise, as to the quality or fitness for any purpose of our goods are excluded, and no responsibility will be accepted by us or resellers for any failure in performance, damage or injury whatsoever arising from their storage, handling, application or use. These conditions cannot be varied by our staff or agents whether or not they supervise or assist in the use of such goods.

**AZOFIN PLUS** is a trademark of Finchimica S.p.A., Italy.