

# Maize Herbicides Product Summary - Spring 2022

Don't get lost in maize weed control choice.

With a choice of leading herbicides providing broad spectrum weed control across a wide range of timings.

Find out more at [www.claytonpp.com](http://www.claytonpp.com)

CLAYTON  
SMELTER®

EC formulation  
960g S-metolachlor

H

CLAYTON  
LAUNCH®

EC formulation  
212.5g dimethenamid-p  
250g pendimethalin

H

CLAYTON  
KIBO®

WG formulation  
750g prosulfuron

H

CLAYTON TASSEL®  
CLAYTON GOLDCOB®

SC formulations  
100g mesotrione

H

CLAYTON NERO®  
CLAYTON BRAMBLE®

WG formulations  
250g rimsulfuron

H

### ***Post-drilling, pre-emergence options for early season weed control in forage and grain maize:***

#### **Clayton Smelter® 960g S-metolachlor EC**

**Use rate & timing:** Apply at 1.4 l/ha post-drilling but pre-emergence of the crop for control of certain annual broad-leaved weeds and grasses.

### ***Pre-emergence and early post-emergence options for weed control:***

#### **Clayton Launch® 212.5g dimethenamid-P, 250g pendimethalin EC**

**Use rate & timing:** Apply at 4.0 l/ha after sowing before crop emergence or post-crop emergence at 1-4 leaves (GS10-14) for control of a range of weeds during germination/early development.

### ***Post-emergence options for weed control:***

#### **Clayton Kibo® 750g prosulfuron WG**

**Use rate & timing:** Apply at 20 g/ha, at 2-4 true leaves stage, for control against a range of annual broad-leaved weeds.

#### **Clayton Goldcob®, Clayton Tassel® 100g mesotrione SC**

**Use rate & timing:** Apply at 0.75 l/ha for susceptible BLWs and 1.5 l/ha for cockspur grass, during 2 and 8 leaves of the crop when weeds actively growing and at seedling stage.

#### **Clayton Bramble®, Clayton Nero® 250g rimsulfuron WG**

**Use rate & timing:** Apply at 50 g/ha, from after crop emergence to before the 4th collar has fully emerged for control of BLWs. Addition of a non-ionic wetting adjuvant at 0.1 % v/v is also recommended.