



OSR Fungicides Product Summary

Maximise your oilseed rape yields this spring.

With an extensive range of reliable fungicides targeting yield damaging diseases.

Find out more at www.claytonpp.com

**CLAYTON
OZARK®**

SC formulation
250g azoxystrobin

F

**CLAYTON
NAVARO®**

EC formulation
125g prothioconazole
125g tebuconazole

F

**CLAYTON
TRELLIS®**

EC formulation
160g prothioconazole
80g tebuconazole

F

**CLAYTON
TUSKAR®**

EC formulation
250g prothioconazole

F

**CLAYTON
TEBUCON® 250EW**

EW formulation
250g tebuconazole

F

METFIN 90™

EC formulation
90g metconazole

F

Clayton Ozark® 250g azoxystrobin SC

Use rate & timing: Apply at 1.0 L/ha for a maximum of 2 treatments per crop at a minimum 21 day spray interval. Latest application 21 days before harvest. Use against *Sclerotinia* as a protectant during flowering (best results from an application timing GS60-GS65, from early to mid-flowering). Use against *Alternaria* as a protectant at early pod formation (before first spots appear on the pods).

Clayton Navaro® 125g prothioconazole, 125g tebuconazole EC

Use rate & timing: Apply at 1.0 l/ha, for a maximum of 2 treatments per crop at a minimum 14 day spray interval. Latest application 56 days before harvest. For Light leaf spot follow up spray(s) may be required in early spring from the onset of stem elongation, depending on disease development. For *Sclerotinia* stem rot, apply at early to full flower.

Clayton Trellis® 160g prothioconazole, 80g tebuconazole EC

Use rate & timing: Apply at 1.0 l/ha, for a maximum of 2 treatments per crop at a minimum 14 day spray interval. Latest application 56 days before harvest. For Light leaf spot follow up spray(s) may be required in early spring from the onset of stem elongation, depending on disease development. For *Sclerotinia* stem rot, apply at early to full flower.

Clayton Tuskar® 250g prothioconazole EC

Use rate & timing: Apply at 0.7 l/ha, for a maximum number of 2 treatments per crop at a minimum 14 day spray interval. Latest application 56 days before harvest. For Light leaf spot, depending on infection pressure, a second spray may be required in spring from stem elongation. For *Sclerotinia* stem rot, apply at first to full flowering.

Clayton Tebucon® 250EW 250g tebuconazole EW

Use rate & timing: Apply at 1.0 L/ha between growth stages BBCH 20 and 69. For Light leaf spot, if disease develops over winter (and no autumn treatment used), apply 1.0 L/ha in early spring from onset of stem extension. Or application either pre- or post-flowering will generally control late development of Light leaf spot on pods and leaves. For *Alternaria* spp., apply 1.0 L/ha at onset of disease when black spots first appear on pods. For *Sclerotinia* stem rot, 1.0 L/ha applied at early to full flower will give some reduction. For Ringspot, spring/summer applications made for Light leaf spot control may give some reduction.

Metfin 90™ 90g metconazole EC

Use rate & timing: Apply at 0.8 L/ha for a maximum of 2 treatments per crop at a minimum 14 day spray interval. Latest application 10% pods at final size. Apply during stem extension, once the crop is actively growing (GS31-51).