

CLAYTON LINAXO
MAPP 20611

GROUP

27

FUNGICIDE

A water dispersible granule containing 600 g/kg cymoxanil for the control of potato blight.

The (COSHH) Control of Substances Hazardous to Health Regulations may apply to the use of this product at work.

Authorisation holder:

Clayton Plant Protection Limited, Bracetown Business Park, Clonee, Dublin 15, Ireland

Telephone: +353-1-8210127 www.claytonpp.com

Contents: 1 to 2kg

FOR PROFESSIONAL USE ONLY

Warning



Harmful if swallowed

May cause an allergic skin reaction

Very toxic to aquatic life with long lasting effects

Avoid breathing spray

Do not eat, drink or smoke when using this product

Wear protective gloves/protective clothing/eye protection/face protection.

IF SWALLOWED: Call a POISON CENTRE or doctor if you feel unwell

IF ON SKIN: Wash with plenty of water:

If skin irritation or rash occurs: Get medical advice/attention

Wash contaminated clothing before reuse.

Call a POISON CENTRE/doctor if you feel unwell.

Collect spillage.

Dispose of contents/container to a licenced hazardous-waste disposal contractor or collection site except for empty clean containers which can be disposed of a non-hazardous waste.

Do not contaminate water with the product or its container (Do not clean application equipment near surface water/Avoid contamination via drains from farmyards and roads).

To avoid risks to human health and the environment, comply with the instructions for use.

SAFETY PRECAUTIONS

OPERATOR PROTECTION

CLAYTON

Engineering control of operator exposure must be used where reasonably practicable in addition to the following personal protective equipment:

OPERATORS must wear suitable protective clothing (coveralls) and suitable protective gloves when handling contaminated surfaces.

However, engineering controls may replace personal protective equipment if a COSHH assessment shows they provide an equal or higher standard of protection.

WASH CONCENTRATE from skin or eyes immediately

DO NOT BREATHE SPRAY

WHEN USING DO NOT EAT, DRINK OR SMOKE

WASH HANDS AND EXPOSED SKIN before meals and after work

IF YOU FEEL UNWELL, seek medical advice immediately (show the label where possible).

ENVIRONMENTAL PROTECTION

EXTREMELY DANGEROUS TO FISH OR OTHER AQUATIC LIFE. Do not contaminate surface waters or ditches with chemical or used container.

STORAGE DISPOSAL

KEEP AWAY FROM FOOD, DRINK AND ANIMAL FEEDING STUFFS

KEEP IN ORIGINAL CONTAINER, tightly closed, in a safe place

KEEP OUT OF REACH OF CHILDREN

EMPTY CONTAINER COMPLETELY and dispose of safely

DO NOT RE-USE CONTAINER for any purpose.

IMPORTANT INFORMATION FOR USE ONLY AS AN AGRICULTURAL FUNGICIDE

Crop	Maximum individual dose (kg/ha)	Maximum number of treatments per crop	Latest time of application
Potato	0.15	8	1 day before harvest

Other specific restrictions

- (1) Non-returnable containers must not be re-used for any purpose.
- (2) A minimum interval of 7 days must be observed between applications.

READ THE LABEL BEFORE USE. USING THIS PRODUCT IN A MANNER THAT IS INCONSISTENT WITH THE LABEL MAY BE AN OFFENCE. FOLLOW THE CODE OF PRACTICE FOR USING PLANT PROTECTION PRODUCTS

DIRECTIONS FOR USE

IMPORTANT: This information is approved as part of the Product Label. All instructions within this section must be read carefully in order to obtain safe and successful use of this product.

RESTRICTIONS

- Allow a minimum of 7 days between applications
- Do not allow packs to become wet during storage
- Do not keep opened packs from one season to another
- Only apply CLAYTON LINAXO in tank-mixture with other fungicides (see COMPATIBILITY).

RESISTANCE

CLAYTON 

When fungicides with the same mode of action are used repeatedly over several years in the same field naturally occurring less sensitive strains may survive, propagate and become dominant in the field. It is therefore advised that whenever possible spray programmes should include fungicides with different modes of action. Please refer to UK Fungicide Action Group (FRAG) guidelines for further details. The combination of multiple active ingredients will also help to reduce the selection and build-up of resistant strains of fungi. Potato fungicides may not give complete protection of blight when the incidence of the disease is severe. A strategy for preventing and managing resistance should be adopted.

The Fungicide Resistance Action Group (FRAG) have produced guidelines and copies are available from AHDB, CPA, your distributor, crop advisor or product manufacturer.

CLAYTON LINAXO contains cymoxanil, a group 27 fungicide, which is locally systemic and offers both preventative and curative control. Apply as part of a season-long programme of correctly timed treatments commencing before blight infection and in tank-mixture with specified products (see COMPATIBILITY).

CROPS

CLAYTON LINAXO can be applied on all varieties of early and maincrop potatoes. On crops grown for processing consult your processor before use.

APPLICATION METHOD

Spray equipment should be thoroughly cleaned before use.

Apply with a tractor mounted/trailed sprayer in 150 litres of water per hectare using a MEDIUM quality spray (as defined by BCPC). Higher water volumes may be used in thick or dense crops to improve coverage

DOSE

Apply CLAYTON LINAXO at 0.15 kg/ha in tank-mixture with another product recommended for control of potato blight containing a fungicide from a different cross resistance group and at a dose rate that will give robust control. All other requirements and restrictions on the partner product label(s) should be adhered to.

TIMINGS

Timing of spray applications should take into account the prevalence of the disease in the area and if conditions support disease spread. The first spray should be applied as soon as there is a risk of blight infection or an official warning has been issued. In the absence of a warning, spraying should begin just before the crop meets along the rows. Applications should be repeated at 10-14 day intervals depending on the prevalence of the disease and favourable conditions for infection. A 7-day spray interval may be used under exceptional circumstances e.g. blight risk is already high and rain is forecast at the time of the next 10-day spray is due. High risk situations include conditions of rapid growth, irrigated crops, areas of intensive potato growing, particularly blight-sensitive varieties, when further warnings are issued, or when active blight is in the area. The longer interval is recommended under dry conditions. Irrigation of crops increases the risk of blight and crops should always be sprayed with CLAYTON LINAXO as soon as possible after irrigation subject to the minimum spray interval. Irrigated crops should normally be sprayed at 10-day intervals

MIXING

Spraying equipment should be clean before use. CLAYTON LINAXO mixes easily with water and should be added to the spray tank during filling. Wash out container and add washings to the spray tank. Continue agitation until spraying is completed.

COMPATIBILITY

CLAYTON

CLAYTON LINAXO is compatible with the following products:

- Fluazinam
- Mancozeb

All other requirements and restrictions on the partner product label(s) should be adhered to.

SPRAYER CLEANING

Immediately after application, completely drain spray equipment. Thoroughly rinse sprayer and flush the hoses, boom nozzles with clean water. Loosen and physically remove visible deposits. Remove and clean nozzles, screens and strainers. Flush the entire system with clean water. Take all necessary safety precautions when cleaning equipment. Do not clean near wells or water sources. Dispose of waste rinse water in accordance with local regulations.

COMPANY ADVISORY INFORMATION

The following information does not form part of the product label under the UK plant protection products regulations and provides additional advice on product use at the discretion of Clayton Plant Protection Ltd.

NOTICE TO BUYER

All goods supplied by us are of a high grade and we believe them to be suitable for any purpose for which we expressly supply them, but as we cannot exercise control over their mixing or use, all conditions and warranties statutory or otherwise, as to the quality or fitness for any purpose of our goods are excluded and no responsibility will be accepted by us for any damage or injury whatsoever arising from their storage, handling, application or use

Since the occurrence of resistance cannot be forecast, neither Clayton Plant Protection Limited nor its distributors can accept responsibility for any loss or damage to crops caused by the failure of CLAYTON LINAXO to control resistant strains.